

Disease and Insect Control Programs for Homegrown Fruit in Kentucky Including Organic Alternatives, 2005

R.E. Durham, J.G. Strang, J.R. Hartman, R. Bessin, R.T. Jones, and J.G. Masabni

Many homeowners in Kentucky grow a variety of fruits in their garden and are rewarded for their effort. One distinct advantage homeowners have over commercial orchardists is the diverse ecosystem of the home landscape (vegetable gardens, flower and fruit plantings intermixed with turf and landscape plants). Diversity often reduces insect and disease organism spread and tends to keep their populations at lower, more manageable, levels.

If the home grower is a careful observer and knows what to look for, pest control can be used as needed, instead of on a rigorous preventive schedule. Arrangements for pest control should be incorporated into plans for any home fruit planting. Failure to understand and carry out insect and disease control measures is an invitation to disaster. On the other hand, use of cultural and biological controls such as sanitation and resistant varieties combined with well-applied sprays of proper materials as needed can bring success in home fruit production.

Cultivar selection, health and vigor of individual plants, managed tree and plant size, good pruning, and weed, grass and rodent control are very important practices for success of home fruit plantings.

Value of Good Cultural Practices in a Pest Management Program

Choose Your Planting Site with Care

Where you plant trees or plants can have a strong influence on future disease problems. Choose a site with good air circulation and sunlight (e.g., away from walls and shade trees) to allow fruits and foliage to dry after periods of rain or dew. Make sure that the soil in your planting site has good drainage. A site where water stands for long periods or collects after a rain is likely to lead to root rot diseases.

Plant What You Need and Can Manage

Avoid too large a planting. A common mistake for new growers is to make the home planting too large, often leading to neglect and discouragement. For a family of four, the following tree and bush numbers are usually sufficient: apples (2); peaches (2); tart cherries (1); plums (2); red or black raspberries (5 plants each of 2 varieties that have different maturity dates); pears (2); grapes (3 vines); thornless erect blackberries (5); thornless semi-erect blackberries (2); blueberries (3); gooseberries (1); and strawberries (25).

Many people find small fruits such as blueberries, strawberries, blackberries, raspberries and gooseberries more rewarding than tree fruits. Small fruits come into bearing sooner, need less space, bloom reliably and require fewer sprays for pest control. Grapes are an exception to this in that they require a rigorous spray schedule and take three years to bear.

Choose Cultivars with Disease Resistance

In general, the home fruit gardener has a much better chance for success when early ripening varieties are used. Early varieties require less spraying and are generally harvested before serious pest problems develop. Use of disease-resistant varieties also helps. For example, several new apple varieties have resistance to some diseases (Table 1).

Keep Your Plants Healthy and Vigorous

Healthy trees and plants yield more fruit than either less vigorous or more vigorous ones. A reasonable supply of mineral nutrients provided by the application of fertilizers and/or organic matter helps assure a harvest. It is important, however, not to apply too much nitrogen to fruit plantings because fruit set is reduced and succulent growth is more susceptible to diseases such as fire blight of apples and pears. A normal fertility program for a lawn usually meets the fertilizer requirements for fruit trees.

Prune Your Plants Correctly

Dwarf or semi-dwarf trees are attractive to home fruit growers because they are easy to spray, prune and harvest. Pruning at planting is very important. Once this is done correctly, prune young trees only enough to shape them until they come into bearing. As trees get older, pruning should be increased.

Prune and thin large trees to improve foliage drying and spray coverage of all parts of the tree. With large trees you may need an extension spray rod and stepladder to get good spray coverage. Remember that with very large trees, the tops can be lowered by heavy limb removal; the remainder can then be sprayed more easily. All sprouts or suckers that arise at the base of trees should be removed.

Late winter or early spring is the best time to prune. Fruit trees should be pruned in the following order to avoid injury from late spring freezes: apples, pears, cherries, plums and peaches. On trees subject to frequent frost loss, pruning may follow bloom and/or fruit set. Limited summer pruning reduces vegetative growth the following spring, so it is a recommended practice on very vigorous trees.

The authors wish to thank Dr. M.L. Witt and Dr. G.R. Brown, UK Professor Emeriti, for their help in preparing this publication.

Variety	Resistance To				Comments	Harvest	Stores Till	Skin Color
	AS	CR	FB	PM				
Pristine**	VR	S	S	S	Good quality for season, not as tart as Lodi, makes excellent applesauce	Early-July	Short Storage	Light yellow with red blush
Williams Pride	VR	S	—	R	Good quality for season, corkspot frequently observed, subacid, yellow flesh	Mid-July	Short Storage	70-80% dark red
Redfree**	VR	VR	S	S	Firm, summer apple, juicy	Late-July	Oct.	90-100% dark red on yellow
Dayton**	VR	S	—	—	Similar to Prima	Mid-Aug.	Sept.	Up to 90% bright medium red
Liberty**	VR	R	R	S	Fruit similar to Macoun, crisp, juicy, yellowish flesh, tart at harvest	Late-Aug.	Dec.	90% dark red stripes on green yellow
Nova Easygro	VR	VR	R	S	Fruit similar to Cortland, fair quality (for trial)	Early-Sept.	Dec.	80% dark red on green yellow
Spartan**	MR	R	MR	R	Firm McIntosh type, needs thinning to develop size	Early-Sept.	Jan.	Dark red to pale red depending on weather
Jonafree**	VR	S	S	R	Fruit similar to Jonathan, but less acid	Early-Sept.	Dec.	90% red stripes
Pixie Crunch**	VR	—	—	—	Small, sweet flavored, super crisp, kids apple	Early-Sept.	Dec.	Deep red
Macfree	VR	MR	MR	S	Similar to McIntosh, mealy under hot conditions	Mid-Sept.	Dec.	75% medium red over green yellow
Priscilla**	VR	MR	VR	R	Tart, firm, somewhat coarse textured, crisp, juicy, small fruit size	Mid-Sept.	Nov.	70-90% dark red blush over yellow green
SirPrize**	VR	S	R	R	Fine grain, crisp, tender, bruises very easily, sterile pollen	Mid-Sept.	Dec.	Greenish yellow slight red blush
CrimsonCrisp	VR	MR	S	S	Medium sized red fruit, firm, crisp, tart, stores very well	Mid-Sept	March	95% red
Enterprise**	VR	VR	—	R	Sprightly, subacid, slightly aromatic and spicy, crisp, fine grained juicy flesh, stores well	Early Oct.	Feb.	Washed, 90% light to medium red
GoldRush**	VR	S	—	S	Fruit very crisp, firm, tart at harvest and sweetens up after storage, very susceptible to black rot. Will store for 11 months.	Mid-Oct.	April	Deep yellow with red blush
Sundance**	VR	VR	VR	VR	Excellent quality with fruity flavor like mild pineapple, fruit does not drop	Mid-Oct.	Mar.	Yellow, occasionally russets in stem cavity

AS = Apple Scab, **CR** = Cedar Apple Rust, **FB** = Fire Blight, **PM** = Powdery Mildew
VR = Very Resistant, **R** = Resistant, **MR** = Moderately Resistant, **S** = Susceptible, **—** = Insufficient Information
Note: All apples require cross-pollination by a different variety. Winesap and SirPrize cannot serve as pollinizers because they have sterile pollen.
* Resistance to diseases other than scab has not been fully evaluated and may differ in some locations from that reported here.
** Produce high quality apples in Kentucky.

Protect Your Trees from Mice or Voles

Field mice of two types often kill many fruit trees. The meadow mouse, a surface worker, gnaws bark from trees at the soil surface or above. The pine mouse tunnels under soil, destroying roots and below-ground trunk. An area clear of vegetation and debris a foot or so around the tree base helps keep mice away. An 18-inch-high cylinder of hardware cloth placed around the tree base and extending 2 inches into the soil gives much better protection. Cats and dogs may help with the mouse problem.

Bird Control

Birds that consume ripening fruit can be a significant problem for home plantings of cherries, raspberries, blackberries, blueberries, currants, gooseberries, strawberries, grapes and sometimes apples, peaches, and plums. Some of the more devastating birds are robins, flickers, red winged blackbirds, crows, cedar waxwings, blue jays, common grackles and European starlings. Federal laws prohibit capture or kill of all species except starlings, house sparrows and feral pigeons. All other birds are protected.

Hanging aluminum pie tins that blow in the breeze, stretching Mylar reflective tape over tops of plants, placing rubber snakes and owls on or above plants, using bird scare balloons with large eyes on their sides and several other techniques all

work to some extent depending on bird populations and amount of other food available in the area at the time. The devices are more effective when employed before the bird problem develops and if they are moved and repositioned frequently. However, birds eventually become accustomed to the scare devices. Thus, they are effective in slowing down losses to birds, but not in eliminating them.

The most effective means of controlling birds is to exclude them with netting. There are several brands of bird netting on the market that may be reused for several years. University of Kentucky researchers have had good success recently on thornless blackberries, grapes and blueberries with several finer-meshed nets that also exclude Japanese beetles and green June beetles.

Insect Traps

Traps can be used to determine when certain pests are present and in what numbers. Traps can also be used to control some pests when the number of trees is small. Traps are often used to monitor the adult form of insects that cause damage in the immature or larval form. Pheromone traps are available for codling moth, peachtree borers, various leafrollers, grape berry moth, oriental fruit moth, dogwood borer, San Jose scale and lesser peachtree borer. These traps use sex-attractant lures to monitor adult males of specific pests.

Pheromone traps can reduce codling moth numbers when many traps (2 to 4 per tree) are used. Pheromone traps need to be set out early in the season, usually before petal fall. Lures in traps should be replaced frequently, as per manufacturer's recommendations. This technique works best when the trees are isolated from other sources of infestation, i.e., other fruit trees. Pheromones are also being sold to prevent mating in some insects such as the codling moth, oriental fruit moth and grape berry moth. Mating disruption requires that a large amount of pheromone be dispensed over an area, and that the orchard is isolated from other sources of these pests. At present, mating disruption is not economically practical for small home plantings. Japanese beetle traps are not a recommended technique for Japanese beetle control. They attract unbelievable numbers of beetles and frequently increase your problem.

Create an Environment Unfavorable for Diseases

Apple and pear scab, powdery mildew, and fruit rot diseases are more serious in moist, shaded locations. These diseases can be reduced by improving air movement through and sunlight penetration into trees. Prune away overhanging vegetation from nearby landscape trees. Thin fruit trees to promote greater sunlight penetration. If watering is needed, irrigate soil around trees rather than using overhead sprinklers, so long periods of leaf wetness are avoided. Plant new trees in well-drained sunny locations. North or east facing slopes make good orchard sites because they have good air drainage and because they are cooler later in the spring, thus bloom is delayed and there is less chance of frost or freeze injury during bloom. If soil is constantly wet, improve soil drainage by ditching and tiling.

Exclude Diseases from the Home Fruit Planting

Some diseases, especially soil-borne and root diseases, may be brought into the planting on transplants. Buy nursery stock from reputable growers. Inspect all purchases for galls, root decay, stem cankers or insect pests and exclude diseased or insect-infested plants from the home fruit planting.

Mites on Fruit Trees

Where mites appear to be a routine problem, the best control strategy is to use several tactics. First, a thorough dormant oil spray at green tip reduces the overwintering population. Second, select insecticides that are least damaging to predatory mites to delay buildup of troublesome populations. Finally, when the number of mites per leaf averages six to 10, apply Vendex and repeat in seven to 10 days. Miticide treatments may not be necessary every year, but when they are, be sure to use a two-spray treatment because most miticides do not kill eggs. Avoid unnecessary use of miticides because it quickly leads to resistance development by mites. If uncontrolled, hatching eggs can quickly lead to a buildup of high populations.

Keep Weeds Out of Your Fruit Plantings

Young trees grow best with clean culture under the tree spread. Older trees can grow and bear well in grass and sod; however, they do best if a bare area is maintained beneath the trees. When present, grass should be mowed, and weeds should not be allowed to grow up through trees. Homeowners should pull and hoe weeds that grow in fruit planting areas. Weedeaters can seriously damage tree trunks, and care must be taken when these are used.

Mulches, applied 2 to 4 inches thick, control weeds effectively. Before choosing a mulch, consider availability, cost, appearance and rodent protection. Mice can hide undetected in mulch; therefore, tree guards are a must. Mulches that are applied once during the growing season cut down on weeds significantly and conserve moisture. Weeds that do come through the mulch are easily pulled by hand. While many preemergence herbicides are labeled for use in commercial fruit plantings, they are not recommended for home fruit plantings.

Control Recommendations and Spray Schedules for Specific Fruit Crops and Pests

The following discussion, grouping similar fruit crops, is intended as a guide for making decisions in controlling specific pests. Sanitation measures to reduce pests are critically important in managing home plantings.

Sprays schedules indicate when specific pesticide applications are required to control various pests on susceptible varieties. Pesticides are used to kill insects and disease pathogens. Handle these chemicals carefully to prevent injury to you, other people, and pets.

If you plan to buy and use chemicals, acquaint yourself with the material to be used, what problems it controls, safety precautions, and specifications of the label. Unless fruit thinning is desired on apples or pears, do not use Sevin in the spray schedule until three to four weeks after flower petals fall. Sevin is especially toxic to honeybees. Repeated use of Sevin increases mite and aphid populations. Always avoid spraying insecticides during bloom in order to preserve honeybees.

The spray schedules are only suggested guidelines because environmental conditions and pest populations vary greatly from year to year. Because of constantly changing pesticide regulations, formulations, and labels, specific application rates are not provided. For more information on common pesticides and the pests they control, refer to the section "Notes on Pesticide Products Commonly Used by Home Fruit Growers" section. It is the applicator's responsibility to obtain the most current information on use directions and application rates directly from the pesticide label.

Apples and Pears

Pest Control Is Aided by Cultural and Sanitation Measures

Apple scab, pear scab and pear leaf spot. All of these diseases overwinter on old dead leaves on the ground. Collect and destroy all leaves in fall, including those from nearby flowering crabapples and ornamental pears.

Cedar apple rust. Spores that infect apple trees come from galls produced by the fungus on cedars. Do not plant susceptible apple varieties near red cedars (junipers). If cedars are part of an established landscape, remove and destroy all galls caused by the rust fungus on cedars in late fall. Inspect cedars again in early spring during or just after a rain when the orange, gelatinous fungus growth from any remaining galls is highly visible.

Fire blight, powdery mildew and summer rots. Prune out and destroy all dead or diseased shoots and limbs while trees are still dormant. Early March is usually a good time to prune trees. All pruning cuts should be made at the base of a branch leaving only the branch collar ($\frac{1}{4}$ to $\frac{1}{2}$ inch) so that

proper wound healing occurs. Removal of dead and broken limbs with cankers and all blighted twigs helps reduce rots and increases effectiveness of sprays when applied. All prunings from trees should be destroyed. As a rule, tree wound paints and dressings should not be used.

Sanitation for fire blight begins by thoroughly examining trees at least once a week from the time blossoms appear in spring until early August. If pruning is to be done during the growing season it must be done just as symptoms appear; **otherwise, it is best to let the disease run its course.** Pruning fire blight out of trees during the growing season creates a risk of increased disease spread. In many cases, pruning can be put off until winter. However, if the disease is serious in young trees and tree structure is at risk, immediate pruning might be warranted. Break or cut infected parts off 6 to 8 inches below the lowest point of visible infection. If removal is made by cuts, avoid spreading bacteria by disinfecting blades of cutting tools between each cut with one of the following, 70 percent alcohol, 10 percent solution of household bleach, Pinesol, or 5 percent Lysol. Before moving on to something else, remove diseased material from the vicinity of the trees and burn, bury, or otherwise dispose of it. Be sure to wash household bleach or Pinesol residue from your tools when finished to prevent rust.

Use disease-resistant cultivars. See Table 1 for a listing of disease-resistant apple cultivars. Avoid apple cultivars such as Jonathan, Yellow Transparent, Gala, Ben Davis, Granny Smith, Jonagold, Melrose, Mutsu, Paulared, York Imperial, Lodi, Idared, Braeburn, Fuji, and Rome Beauty, which are very susceptible to fire blight. Pear cultivars resistant to fire blight include Magness, Maxine, Seckel, Moonglow, Potomac, Harrow Sweet and HoneySweet. Bartlett, Comice, Anjou, Clapps Favorite and Bosc pears are very susceptible to fire blight and should not be planted.

Use nitrogen fertilizers very sparingly around fire blight-susceptible pear and apple cultivars. Succulent growth that results from excessive nitrogen application encourages development of the disease once inoculated.

Apple maggot, codling moth and plum curculio. Fruits on the tree showing entrance or exit wounds (holes) of either apple maggot or codling moth should be removed and destroyed immediately. Apple maggot is rarely found in Kentucky, but is a serious problem north of the Ohio River. Similarly, all fruits that fall prematurely, especially during May and June, should be promptly collected and destroyed. This is an important step because it prevents developing larvae from reaching maturity and causing more damage to other fruits later in the season. Codling moth larvae are attracted to corrugated cardboard strips as a place to pupate over winter. Attach strips of corrugated cardboard around the trunk of the tree in late summer and remove and destroy them in winter.

Pear Psylla on Pears

Sometime just before the green tip stage of pear bud development, tiny pear psylla nymphs appear and begin sucking plant juices. They collect beneath leaves and if left unchecked, their feeding in summer causes leaves to turn brown and drop. Leaves, twigs, and fruits are often covered with sticky honeydew (psylla droppings) on which a black, sooty mold grows.

If pear psylla has been a problem, apply dormant oil spray before buds open, followed by an insecticide at petal-fall and in the first cover spray. Wait 30 days and if psyllids are still present, make two more applications of insecticide seven to 10 days apart.

Pesticide Spray Program

A preventive approach emphasizing early-season control of all major diseases and a number of important insect pests is recommended (Table 2). The critical time period is between bud break (green tip) and about one week after petal fall. This is the only time that certain diseases (scab, apple rust and powdery mildew) and insect pests (plant bugs) can be controlled effectively. In other cases (where pear leaf spot, mites, aphids, plum curculio, pear psylla and leafrollers are problems), a good early-season program not only provides excellent control, but also greatly reduces potential for late-season damage and, therefore, the need for frequent sprays later in the season. Where an early-season control program is followed, timing of mid- to late-season sprays is much less critical and the interval between applications can be safely lengthened to two or three weeks. Learning to identify insect and disease symptoms is the key to reducing sprays at this time.

Peaches, Nectarines, Apricots, Plums and Cherries

Brown rot is the most important disease affecting all stone fruit (peaches, nectarines, plums, and sweet and tart cherries). However, other diseases such as peach leaf curl, *Cytospora* canker, plum black knot and cherry leaf spot also should be considered. Borers, catfacing insects (plant bugs, stink bugs) and the oriental fruit moth are usually the most destructive among insect pests that attack peaches. Peachtree borer also attacks other stone fruit trees. Plum curculio can destroy an entire plum crop and is harmful to other stone fruits, as well (Table 3.)

Sanitation Measures for Disease Control

Brown rot. Brown rot fungus overwinters on old mummified fruit left hanging on the tree or on the ground. After harvest or in spring before buds open, collect and destroy all mummified fruits beneath trees and any that remain hanging in trees. Collect and destroy any fruit that falls prematurely or shows symptoms of brown rot during the growing season. Never allow rotting fruit to remain on or under trees. Do not allow fruit to over-ripen or soften before harvesting. Peaches, nectarines, and apricots should be harvested when their color (nonblush side) first changes from green to all yellow, even though the flesh is still firm. Harvest plums when fruit color is uniform and flesh begins to soften under gentle pressure. Two to three days of further ripening off the tree are necessary to achieve better eating quality. However, tree-ripened fruit have the best eating quality. Good air circulation through the tree is essential to reduce wetness, which favors brown rot. Pruning allows better air and sunlight penetration and reduces disease incidence.

In Kentucky, winter injury, spring frosts during bloom, and *Cytospora* canker are major problems for stone fruit. Consequently, homeowners should wait until after bloom to prune stone fruits when it is easier to determine the few flowers

Time to Spray	Examples of Pesticides Required ¹			Most likely insect and disease problems at this time.
	Apple	Pear	Organic Pesticides	
Dormant Early spring before buds swell	Dormant oil Bactericide: fixed copper Note: Do not spray dormant oil when temperature is below 40°F (5°C), or likely to drop below 40°F within 24 hours.	Dormant oil	Dormant oil Bactericide: fixed copper	Oil for scale. Check label carefully for dormant vs. delayed dormant rates on apple vs. pear. Bactericide for fire blight. Do not mix bactericide with oil. Copper sulfate is not the same as fixed copper, and should not be mixed with oil.
Green Tip to half-inch green When blossom buds show ½ inch green	Fungicide: captan*, mancozeb, ferbam, thiophanate-methyl + captan*, or myclobutanil	No spray needed	Fungicide: fixed copper or sulfur	Fungicide for control of scab. Last chance to apply oil on apples, if not applied in dormant spray. Use dormant oil now instead of at dormant if scale is not a problem.
Tight to open cluster When fruit buds are visible	Fungicide: same as green tip	No spray needed	Fungicide: same as green tip	Fungicide for control of scab. Using fixed copper past this point is likely to cause fruit russetting.
Pink Just before blooms open	Fungicide: same as green tip plus ferbam, mancozeb + Insecticide: Thiodan or Malathion	Fungicide: see apple	Fungicide: same as green tip + Insecticide: Sabadilla or Pyrethrum	Fungicide for control of scab and rust. Insecticide for aphids, tarnished plant bug and stink bug if present.
Bloom When 20%-60% blossoms are open	Fungicide: same as green tip plus bactericide streptomycin if available and appropriate.	No spray needed except bactericide if needed: See apple.	Fungicide: fixed copper	Fungicide for scab, rust and powdery mildew. Bactericide for fire blight. Note: do not use insecticide during bloom.
Petal fall When last petals are falling	Fungicide: same as pink + Insecticide: Malathion or Thiodan	Fungicide: see apple + Insecticide: see apple	Fungicide: same as green tip + Insecticide: same as pink	Fungicide for control of scab, rust, fruit rots and sooty blotch. Insecticide for codling moth, plum curculio, pear psylla, plant bugs and leafrollers.
First Cover Seven days after petal fall spray	Fungicide: captan*, or thiophanate-methyl + captan* + Insecticide: same as petal fall	Fungicide: see apple + Insecticide: see apple	Fungicide: sulfur + Insecticide: same as pink	Fungicide for control of scab, rust, fruit rot and sooty blotch. Insecticide for codling moth and plum curculio.
Second Cover Two weeks after first cover spray	Fungicide: same as first cover + Insecticide: same as petal fall + Miticide: Vendex	Fungicide: see apple + Insecticide: see apple	Fungicide: same as first cover + Insecticide: same as pink	Fungicide for control of scab, rust, fruit rots, sooty blotch and leaf spots. Insecticide for codling moth, plum curculio, San Jose scale crawlers. Miticide for mites only, if present.
Third Cover Two weeks after second cover	Fungicide: same as first cover + Insecticide: same as petal fall	Fungicide: see apple + Insecticide: see apple	Fungicide: same as first cover + Insecticide: same as pink	Fungicide for control of scab, fruit rots and sooty blotch. Insecticides for codling moth and pear psylla.
Remaining covers Spray every two weeks	Fungicide: same as first cover** + Insecticide: same as petal fall	Fungicide:** see apple + Insecticide: see apple	Fungicide: same as first cover + Insecticide: same as pink	Fungicide for control of fruit rots, scab and sooty blotch. Insecticides for codling moth, leafhoppers and pear psylla. Always observe all pre-harvest intervals.

¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting. See Table 9 for information on mixing small quantities of pesticides.

* Although cleared for use on apples, captan is not labelled for pears.

** Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.

that survive in a year that has a severe winter or spring frost. An additional reason for delaying pruning is to reduce the incidence of Cytospora canker.

Cytospora canker. Be alert for signs of Cytospora cankers. These cankers are dark, sunken areas on limbs and shoots with gum protruding through the bark, often with a raised callus margin. The canker pathogen often enters major limbs through old wounds, broken limbs, and weak shoots killed by low winter temperatures.

Delay pruning stone fruit trees until late April. Pruning at this time helps reduce Cytospora canker because wounds heal quickly.

Pruning weak water sprouts in the center of the tree during late summer (early August) helps prevent establishment of cankers within the main tree structure. Small cankers on shoots and limb tips should be removed completely during the pruning operation. Little can be done once these cankers grow larger or become established in the crotch or on the trunk. Do not use commercial wound paints on pruning cuts. Make cuts so that only a small, raised branch collar remains on the supporting branch, and allow them to close naturally. Limb damage that occurs during the growing season should be repaired immediately; never wait for the dormant season.

Time to Spray	Examples of Pesticides Required¹				Most likely insect and disease problems at this time.
	Peaches Nectarines Apricots	Plums	Cherries	Organic Pesticides	
Dormant for peaches nectarines & apricots Early spring before buds swell or in fall after leaf fall.	Fungicide: ferbam, Bordeaux or chlorothalonil (for leaf curl)	Fungicide: ferbam, Bordeaux or chlorothalonil (only if plum pockets are a problem)	No spray needed	Fungicide: Bordeaux for leaf curl or plum pockets	This is the only time that a fungicide spray will control peach leaf curl and plum pockets.
Bud swell just before the buds break open in the spring	Oil	Oil	Oil	Oil	Oil for control of scale insects, European red mite and aphids. Application is not required if these pests have not been a problem.
	Note: Do not spray oil when temperature is below 40°F (5°C), or likely to drop below 40°F within 24 hours.				
Pink Just before blooms open	Insecticide: Thiodan or Sevin	Fungicide: liquid lime sulfur, thiophanate-methyl, or chlorothalonil	No spray needed	Fungicide: liquid lime sulfur + Insecticide Pyrethrum (peaches)	Insecticide for control of catfacing insects (plant bugs and stink bugs) and early plum curculio. Fungicide is required on plums only if black knot is a problem. For black knot control, fungicides will not be effective unless all knots are removed from the tree and destroyed. Infections on nearby trees must also be eradicated.
Bloom	Note: To protect bees do not use insecticide during bloom.				Fungicide during bloom should not be required if good sanitation is used to control brown rot.
Petal fall When last petals are falling	Fungicide: sulfur, captan, chlorothalonil, myclobutanil, or thiophanate-methyl + Insecticide: Thiodan*	Fungicide: sulfur, captan, chlorothalonil, myclobutanil, or thiophanate-methyl + Insecticide: Thiodan*	Fungicide: captan, chlorothalonil, myclobutanil, or thiophanate-methyl + Insecticide: Thiodan*	Fungicide: sulfur + Insecticide: Same as pink or Surround	Insecticide for control of plum curculio, oriental fruit moth, plant bugs and stink bugs. Fungicide for control of scab and brown rot on all fruits. Also for black knot of plum and cherry leaf spot.
First cover 7 days after petal fall spray	Same as petal fall	Same as petal fall	Same as petal fall	Fungicide: sulfur + Insecticide: Same as pink	Insecticide for control of same pests as above. Fungicide for control of scab, brown rot, cherry leaf spot and black knot of plum.
Remaining covers Continue spraying at 10-to 14-day intervals.	Same as petal fall	Same as petal fall	Same as petal fall. After this spray no further spray needed.	Fungicide: sulfur + Insecticide: Same as pink	Insecticide for control of same pests as above. Fungicide for scab, brown rot, cherry leaf spot and black knot of plum. Use shorter interval for fungicide if wet, rainy weather persists.
Final spray Within one week of harvest	Fungicide**: sulfur, captan, thiophanate-methyl, or myclobutanil	Fungicide**: sulfur, captan, myclobutanil, or thiophanate-methyl	No spray needed	Fungicide: sulfur	For control of brown rot.

¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting. See Table 9 for information on mixing small quantities of pesticides.
** Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.
* Thiodan may only be applied twice in a single season to peaches, plums, or cherries.

Black knot of plum. On plums, in addition to Cytospora cankers, look for symptoms of black knot disease at the time of pruning. These are rough, black tumors or overgrowths that develop on shoots and limbs. They should be pruned out completely. Remove knots from the orchard and destroy them. If possible, also remove wild plums and wild cherries near the planting.

Cherry leaf spot. Cherry leaf spot fungus overwinters on fallen dead leaves. Rake the area under the tree thoroughly and destroy all old leaves to control cherry leaf spot disease. Do this in the fall as soon as leaves drop and in early spring,

before bloom. This procedure can be very effective for small plantings isolated from other cherries.

Plum curculio. Promptly collect and destroy any fruit that falls prematurely to reduce future plum curculio populations.

Yellowjackets, bees, wasps. Plan on harvesting stone fruit over a period of one to two weeks. Regularly pick fully ripened fruits, and remove any that have fallen or any showing decay. This discourages yellowjacket, bee, and wasp populations around and in your trees so that no special insecticide treatments are necessary. It is a good idea to inspect every fruit before grasping it.

Peachtree Borers

The peachtree borer works beneath the bark at or near ground level. The lesser peachtree borer may be found as high as 3 feet above the ground. Presence of borers is indicated by masses of gum that contain small brown particles of bark at the base of the tree, or occurrence of frass and empty pupal cases protruding from tree wounds. If borers are found, they should be killed using a knife or flexible wire to probe larva from the trunk. This should be done carefully; carelessness may result in more damage to the tree than the damage that would have been caused by the borers! See ENTFACT-200, Peachtree borer, for more information. Adult borers begin emerging in late April to mid-May in southern areas, and late May to early June in more northern areas. The lesser peachtree borer is a similar pest that attacks trees anywhere in the lower few feet of the trunk and scaffold limbs rather than at ground level.

It is important to prevent borer damage by protecting the base or trunk of the tree with an insecticide. Insecticide solutions are also effective when applied with a paint brush to the affected area on the lower trunk. Apply Thiodan or Lindane between mid-May and early June and again six to eight weeks later. Do not apply these compounds to the fruit or foliage. Observe the days to harvest restrictions for these insecticides. Avoid mechanical injury to the trunk, since open wounds attract borers. Pheromone traps can be used to detect when moth flights occur so insecticide applications can be timed to coincide with egg laying on trees.

Pheromone traps are available for both peachtree borer and lesser peachtree borer.

Raspberries and Blackberries

Most pesticide treatments on blackberries and raspberries can be made on an “as needed” basis (Table 4). However, this approach requires regular inspection so that needed applications can be made before serious damage occurs. In addition, a good sanitation program in and around the planting helps alleviate a number of common pest problems.

Sanitation Measures

Most fungi that infect raspberry and blackberry canes overwinter on old canes that were infected the previous season. After harvest, cut and remove canes from the planting that have fruited (except for Heritage or other fall-fruited raspberry cultivars) to reduce the incidence of spur blight, cane blight and anthracnose, as well as populations of cane and crown borer insects. Mow or cut old fall fruiting raspberry canes in early spring before new shoots begin to develop. Cuts should be made at soil level (no stubs), and all old canes should be removed from the planting. Spring removal results in production of a fall crop only.

Good weed control promotes rapid drying conditions within the fruiting canopy and discourages populations of insects and diseases that damage canes and fruit. Also, to promote rapid drying, do not allow the row width to exceed 2 feet.

Each spring when new shoots are about 12 to 14 inches tall, survey black raspberry and blackberry plantings for symptoms of orange rust and viruses. Orange rust, a destructive disease of black raspberry and blackberry, can be controlled only by removing all infected plants (including their roots) when symptoms first appear. Infected shoots are identified by their thin, willowy growth, absence of thorns (on normally thorny varieties), and presence of orange spore pustules on the underside of leaves. Failure to remove orange rust-infected plants threatens survival of nearby healthy plants. Virus-infected plants may have leaves with yellow splotches or lines and crumbly, incomplete fruit. Virus-infected plants must also be removed from the planting.

Problems with sap beetles, wasps, fruit flies and fruit rots are less severe if berries are harvested regularly throughout the ripening period so that overripe fruits do not accumulate. Damaged or fermenting fruit attracts these pests.

Japanese beetles feed on ripe raspberry and blackberry fruit and leaves, as well as on fall raspberry flowers. They can be a serious problem in areas with recently established Japanese beetle populations. The spray schedule for brambles is listed in Table 4.

Time to Spray	Examples of Pesticides Required¹	Organic Pesticides¹	Most likely insect and disease problems at this time.
Bud break When buds begin to break and show silver	Fungicide: (Liquid lime-sulfur preferred)	Fungicide: Liquid lime-sulfur	Lime-sulfur for rose scale, anthracnose, cane blight and spur blight. This spray is essential for good disease control. If applied to green tissue, damage may occur.
During bloom	Fungicide: sulfur	Fungicide: sulfur	Fungicide for control of Botrytis fruit rot. In dry growing seasons, fungicide is generally not required. Note: No insecticide used during bloom to protect bees.
Pre-harvest As fruit begin to color	Insecticide: Sevin, Malathion**		Insecticide for sap beetle, cane borer and Japanese beetle, as needed.
Post-harvest	Insecticide as needed. Sevin or Malathion		Insecticide for Japanese beetle and cane borer, as needed.

¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting. See Table 9 for information on mixing small quantities of pesticides.
** Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.

Strawberries

Mites on Strawberries

Mite damage appears as distorted and discolored leaves and stunted fruits. Silken webs may also be found on the lower surface of the leaves. Two species of mites found on strawberries are the two-spotted spider mite, which varies in color from pale greenish yellow to dark crimson with two dark spots on its back, and the cyclamen mite, which is so tiny that it is scarcely visible to the naked eye. Both mites suck sap from foliage, and heavy populations can cause serious damage to strawberry plantings. Life cycles of both species are quite short during summer, and under ideal weather conditions they can build up rapidly if left uncontrolled. Poor weed control can lead to early mite buildup.

When mites are a problem, growers should use a miticide. For best mite control, apply two sprays of insecticidal soap spaced three to five days apart. Hot, dry weather favors rapid development of the two-spotted spider mite, particularly in July and August. Control measures should be applied if mite populations reach five per leaflet. Populations of cyclamen mite usually begin to increase in early May when blossoming starts and peak in early June during fruiting. A sharp decline in cyclamen mite populations generally occurs during July and August. Observations for cyclamen mite probably are best made before blossoms appear. Control measures should

be applied at the first sign of an infestation.

Sanitation Measures

Good weed control is by far the most important sanitation measure for strawberry production. Weeds compete strongly with shallow-rooted strawberry plants and interfere with good runner-plant production. Weeds also harbor many insect and mite pests and prevent rapid drying after rains, promoting increased fruit rot. A good straw mulch in place before fruit begins to ripen helps reduce fruit rot by keeping berries from touching or being splashed with soil.

To reduce populations of sap beetles, remove overripe, diseased and damaged fruit at regular intervals.

Renovate beds immediately after harvest to reduce pest problems. Renovation involves narrowing rows to remove older non-productive plants, weed removal, leaf mowing if foliar diseases are a problem, and fertilization to promote development of new daughter plants that will produce next year's crop. Rake and destroy cut-off leaves and stems after renovation. Select and use varieties with resistance to leaf diseases, Verticillium wilt and red stele.

When old plantings are replaced by new ones, select a different growing site to prevent buildup of soilborne diseases, insects and weeds. The strawberry spray schedule is in Table 5.

Time to Spray	Examples of Pesticides Required¹	Organic Pesticides¹	Most likely insect and disease problems at this time.
Pre-bloom When blossom stems have pushed out of the crown.	Insecticide: Thiodan or Sevin Miticide: Insecticidal soap	Insecticide: Pyrethrum Miticide: Insecticidal soap	Insecticide for aphids, weevil, spittlebug, plant bugs, bud clipper and crown borer. Miticide for spider mites. Note: Early season applications are required only if these pests are a problem.
During bloom	Fungicide: captan NOTE: No insecticide during bloom to protect bees.		Fungicide for fruit rots and leaf spots. In dry growing seasons, fungicide is generally not required.
Post-bloom Begin 10 days after full bloom spray (if needed) and continue spray every seven days as needed.	Insecticide: Thiodan or Sevin		Insecticide for spittle bug, bud clipper and tarnished plant bug. Follow label instructions and observe all pre-harvest intervals.
Harvest Post-harvest Apply one or more times after renovation (if needed) to protect the new foliage for next year's crop.	Fungicide: captan (Use only if needed.)	Diatomaceous earth	Slugs Fungicide for leaf spot

¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting. See Table 9 for information on mixing small quantities of pesticides.
** Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.

Grapes

Sanitation Measures

Black rot is the most common disease home gardeners encounter when producing grapes. Berries infected by black rot fungus turn into dry, shriveled, raisin-like "mummies." The black rot fungus overwinters primarily on these mummies, but also on infected canes. Therefore, it is essential to remove all clusters of mummified fruit from the planting, as well as those that have fallen to the ground. This is best done at harvest or soon after, while they are still easily visible. It is also important to destroy or remove infected canes that are pruned off during the dormant season.

If black rot lesions appear on leaves before fruit set, removing infected leaves should reduce the level of infection.

Most fungi require water on plant surfaces for infection. To promote rapid drying conditions within the foliar canopy, keep vines well pruned (during the late dormant season), trellised and weed-free.

Several pickings are necessary during harvest because not all grape bunches ripen at the same time. Harvesting in this manner reduces incidence of yellow jackets, bees and wasps feeding on overripe fruits, and discourages rot organisms from becoming established. Remove all old, dried or rotted unharvested grapes (including cluster stems) from the vine and destroy them. Collecting and destroying leaf debris under vines in the fall reduces overwintering pupae of the grape berry moth. A spray schedule for grapes is shown in Table 6.

Time to Spray	Examples of Pesticides Required ¹	Organic Pesticides¹	Most likely insect and disease problems at this time.
Dormant	Fungicide: fixed copper or lime sulfur	Fungicide: fixed copper or lime sulfur	Fungicide to control anthracnose, dead arm and black rot.
Bud swell, before buds show green	Insecticide: Sevin	Insecticide: Pyrethrum	Insecticide may be needed if flea beetles or climbing cutworms are a problem (e.g., swollen buds have holes or are eaten). If neither pest is known to occur regularly, no spray is needed.
New growth 4 to 6 inches long	Fungicide: ferbam, mancozeb or captan	Fungicide: fixed copper	Fungicides for control of black rot and Phomopsis cane and leaf spot.
New growth 10 to 15 inches long or 7 to 10 days after first spray	Fungicide: ferbam, mancozeb, captan, or myclobutanil	Fungicide: fixed copper	Same as above
Pre-bloom first blossoms open	Fungicide: ferbam, mancozeb, captan, sulfur, or myclobutanil	Fungicide: fixed copper or sulfur	Fungicide for control of black rot, powdery and downy mildews.
Bloom	Fungicide: fixed copper or sulfur Insecticide: Thiodan	Fungicide: fixed copper or sulfur	Insecticide for control of grape phylloxera. See label; several varieties are very sensitive to Thiodan injury.
Post-bloom blossoms have fallen	Fungicide: ferbam, mancozeb, captan, sulfur, or myclobutanil Insecticide: Sevin or Malathion Miticide: Insecticidal soap Note: In wet weather, do not wait until all blossoms have fallen, especially if black rot is a problem. Spray fungicide every 7 to 14 days.	Fungicide: fixed copper or sulfur Insecticide: Bt	Fungicide for black rot, downy mildew and powdery mildew. Insecticides for grape berry moth and leafhoppers. Miticide for mites if present.
First cover 7 to 10 days after post-bloom	Fungicide: ferbam, captan, sulfur, or myclobutanil Insecticide: same as post-bloom Miticide: same as post-bloom	Fungicide: fixed copper or sulfur Insecticide: insecticidal soap, Bt	Fungicides for control of black rot, powdery mildew and downy mildew. Insecticides for control of grape berry moth, leafhoppers, and Japanese beetle. Miticide for mites if present.
Second cover 10 to 14 days after first cover	Same as first cover	Same as first cover	Same as first cover
Third cover 10 to 14 days after second cover	Same as first cover	Fungicide: fixed copper or sulfur Insecticide: Bt	Fungicide for control of black rot, powdery mildew and downy mildew. Insecticide for grape berry moth and Japanese beetle.
Fourth cover 10 to 14 days after third cover	Same as first cover Always observe pre-harvest intervals.**	Same as third cover	Same as third cover
Special Notes: Berries are no longer susceptible to black rot when they reach about 6 percent to 8 percent sugar content (usually when they start to change color).			
¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting. See Table 9 for information on mixing small quantities of pesticides.			
** Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.			

Blueberries

As a rule, blueberries require the least amount of pesticide of the fruit crops in this publication (Table 7). The best approach is to apply no pesticides to new plantings until some problem develops. Watch plants closely for any symptoms of disease (die back or fruit rot) or insect damage, and begin spraying when appropriate. Some fruit loss can be expected the first year that mummy berry disease or fruit rot develops. In subsequent years, this can be controlled with fungicides.

Probably the most common problems associated with blueberries are birds, nutrient deficiencies related to soil pH requirements and water imbalance from lack of proper mulching and irrigation.

Sanitation Measures

Before buds break in the spring, thoroughly rake the area beneath and immediately around plants to collect or bury any mummified fruits from the previous year's crop. This helps reduce incidence of mummy berry disease. Avoid excessive nitrogen fertilization that increases plant susceptibility to Botrytis fungus (blight). To reduce fungal canker diseases, prune out and destroy dead twigs, branches and stems.

Time to Spray	Examples of Pesticides Required¹	Organic Pesticides¹	Most likely insect and disease problems at this time.
Dormant Before bud break	Dormant oil	Dormant oil	Apply only if scale insects are a problem.
Petal fall 75% petals have dropped	Insecticide: Malathion	Bt	Insecticide for leafroller and plum curculio if present. Bt will not control plum curculio
After bloom 7 to 10 days after petal fall	Same as above Observe all pre-harvest intervals**		Same as above, plus blueberry maggot and Japanese beetle if a problem. Use only if needed.

¹ Growers must read the pesticide label for proper rates of chemical to use. Some materials are effective against some pests and not others. Choose materials needed to control the most important pests in your fruit planting.

** See Table 9 for information on mixing small quantities of pesticides. Check label for waiting days to harvest. Tables 10 and 11 also give waiting days to harvest.

Protecting Yourself When Using Pesticides

Pesticides are used to kill insects and disease pathogens. Handle these chemicals carefully to prevent injury to yourself, other people, or pets. Although pesticides suggested for use in this bulletin are the least hazardous available, certain precautions are still necessary.

1. Before purchasing, mixing, storing, disposing of, or using any pesticide, carefully read the label. Give special attention to the section on various precautions to be followed. Mix only as much as you need.
2. Wear long-sleeved shirts, long pants, rubber gloves, and a hat when mixing and applying chemicals. Goggles should be worn, especially if you wear contact lenses. The product label specifies required minimum protective gear.
3. Avoid spilling pesticides on yourself or in the immediate area where you are working. If this happens, wash yourself immediately with plenty of water to remove all traces of pesticides. Do not get any pesticide in your eyes, nose, or mouth.
4. Do not smoke, eat, or chew tobacco while you are applying pesticides.
5. When applying a pesticide, do not permit material to blow back on you or on other people, pets, or pet food or water containers. Adjust your treatment according to wind direction. If it is too windy, stop and finish later when the wind dies down. A good time to make pesticide applications is in the evening just before dark when the wind usually dies down. Remember, you are responsible for pesticide spray drift.
6. Thoroughly wash yourself and your clothes immediately after applying pesticides.
7. Do not permit empty pesticide containers to lie around; triple rinse containers, wrap in paper, and put them in the trash can. Punch holes in empty containers. Do not burn such containers in a backyard trash burner.
8. Store pesticides in tightly closed, well labeled, original containers away from children or pets, never under the sink, in the pantry, or medicine cabinet. Store in a cool, dry place. Placing bags of wettable powders in a sealed clear plastic bag reduces unpleasant fumes. Mark the storage cabinet or storage area "POISON STORAGE." It is best to keep pesticides under lock and key.
9. Keep children and pets away from areas where you are mixing or applying pesticides.


10. Do not store diluted spray mixtures in jugs or the spray tank from one spray application to the next. It will not be effective and is unsafe.
11. Observe all harvest intervals and reentry requirements after using the pesticide.
12. Keep a written record of what you apply.

Pesticide Training Available

Pesticide training is available at your local county Extension office. If you feel uncomfortable about handling pesticides and understanding labels, ask about the availability of private applicator pesticide certification training.

Spray Equipment for Home Fruit Growers

Thoroughly cover fruit plants with pesticide sprays according to the "timing" in the spray schedule to adequately control pests. Use a sprayer that is powerful enough to reach all parts of the plant with spray (e.g., tops of trees), easy to clean, and slow to wear out. Suitable sprayers come in various shapes and sizes. Refer to Table 8 in deciding what size sprayer is needed for your fruit planting.



Hand pump sprayers like the one pictured here have metal or plastic tanks which vary in size from one to three gallons. Air is pumped into them by a built-in hand pump. Spray is delivered through an attached hose with a hand shut-off valve and nozzle tip.

Hand pump backpack sprayers are compressed air sprayers that vary in size from three to five gallons and are strapped onto the applicator's back. These sprayers, also equipped with hand shut-off valves, have a hand pump that must be pumped slowly but continuously. Pumping builds up pressure in the tank and forces spray through a hose and nozzle

tip at an even, steady rate.

Hand-held compressed air and knapsack sprayers are satisfactory for a few dwarf fruit trees, vines, bushes or strawberry plants. They do not have the capacity to spray mature standard-sized trees.

Trombone or slide-type sprayers consist basically of two small-diameter tubes. One tube slides within the other, compresses liquids and forces pesticide solution through a small

Table 8. Approximate Amount of Spray Required for Fruit Trees of Different Sizes

Height (in feet)	Spread (in feet)	Gallon per tree per application*
4	3	½
5 to 8	3 to 6	1 to 1 ½
8 to 10	4 to 8	2 to 3
10 to 15	8 to 15	3 to 6
15 to 20	15 to 25	5 to 10

* Use the greater amounts for trees in full foliage.

hole in the end of one tube. These sprayers can deliver spray to tops of most fruit trees and are suitable for plantings of a few trees, as well as small fruits.

Small power sprayers such as **backpack mist blowers**, gas power sprayers and electric power sprayers that run off of a 12-volt battery are available for larger home orchard plantings.

Accessory Equipment

Two other pieces of equipment are beneficial to the home fruit grower: a one-quart graduated measuring cup and a set of measuring spoons. **Mark the measuring cup and spoons with spray paint or an indelible marker so that they are used for pesticide measurements only. Keep the cup and measuring spoons separate from those used in the home and store them with the pesticides.** These pieces are necessary to accurately measure required amounts of pesticides, thus ensuring the best control and the least possibility of plant injury. A container designated for making a slurry of the pesticides with water will help in getting the pesticide solution into the spray tank.

How to Spray

Keys to successful pest control are (a) thorough and proper coverage, (b) correct timing, (c) proper dosage, and (d) use of correct materials. These points are extremely critical. Lack of attention to them accounts for poor quality fruit production. In addition, keep the following points in mind:

- Thorough coverage of all above-ground plant parts is necessary to control fruit insects and diseases.
- Direct spray onto both top and bottom sides of leaves until spray begins to drip off the leaves.
- The goal of spraying is to place a thin layer of pesticide over all exposed surfaces of each plant so that no matter where an insect may eat or crawl on that plant or where a disease organism may be, each is exposed to a lethal amount of pesticide before damaging the plant.
- Fungicides prevent infection; therefore, new growth must be protected. Fungicides should be applied more frequently in spring when plants are growing rapidly and when pathogens are active.
- If pest control results are poor after using the spray schedule, be sure to check these important points:
 - Did plants receive thorough spray coverage at the recommended dosage?
 - Were any recommended sprays omitted?
 - Was the timing according to schedule?
 - Did rain occur before the pesticide had time to dry on foliage?
 - Was a fresh spray mixture used each time? Pesticide solutions left in the sprayer soon lose their toxicity and may harm sprayer parts.

Wettable powder sprays tend to settle to the bottom of the tank unless the sprayer has a mechanical or automatic agitator. Stir or shake mixture often during application.

Re-Entry After Spraying

Home orchards are exempt from worker protection standards, which are mandatory for commercial fruit growers. All pesticides have a minimum 12- to 96-hour re-entry interval after spraying, which must be adhered to by commercial growers. **We suggest that you stay out of pesticide-treated areas for at least 12 hours and until the area is dry.**

Cleaning Spray Equipment

Spray any excess spray on the fruit planting rather than dumping it out and contaminating the environment. Next, place a small amount of water in the tank, agitate it and spray this out on the ground beneath the fruit plants. Then, flush the tank several times with clean water and force water through the spray wand and nozzle until clear water is expelled. To prevent corrosion, suspend tank upside down with lid removed to permit drainage and drying.

How Long Pesticides Last

Many pesticides volatilize or lose effectiveness in a matter of days or weeks after being applied. Factors such as temperature, humidity, wind, and sunlight affect the life of pesticides. The greater the extremes of these factors, the quicker pesticides lose their toxicity.

Rain, to some degree, physically removes pesticides from plant foliage. In general, a pesticide is less likely to be washed off if it has had an opportunity to dry thoroughly on foliage before rain. Most materials should be reapplied the day after a heavy rain. Strong sunlight and driving winds also shorten the effective life of pesticides.

New plant growth early in the season results in unsprayed and unprotected parts if spray applications are not repeated at regular intervals.

The above items and the fact that pest populations are continuously moving and/or multiplying are reasons why repeated spray applications are needed.

Storage Life of Pesticides

Although most pesticides, when properly stored, retain their toxicity for use a second or third season, it is best to buy only enough for one season's use. Most pesticides gradually lose their effectiveness when exposed to moisture, air, light and high temperatures. Prolonged low or freezing temperatures frequently cause liquid formulations to separate, making them unsafe and non-effective for further use.

Spray Dilution Chart for Small Quantities of Pesticides

Table 9 is provided as a guide for home fruit growers to determine how much pesticide is needed to mix up small amounts of spray material, because some pesticide containers do not provide this information. **This table should not be used as a recommendation of pesticide rates to apply to fruit crops, but as a guide for mixing pesticides based on the pesticide container label.** There are a number of reasons for this.

One of the major limitations in converting from pounds per 100 gallons to a volume measure such as tablespoons is that wettable power pesticides can have different densities. For

example, Thiophanate-methyl is quite light in comparison to Captan. Thus, there is quite a difference in the volume occupied by a pound of Thiophanate-methyl and a pound of Captan.

Be further cautioned that the rate of a pesticide that is applied may vary between different fruit crops. **Thus, ALWAYS USE THE PESTICIDE LABEL AS THE BASIS FOR YOUR CALCULATIONS.**

Table 9. Spray Dilution Chart for Small Quantities of Pesticides		
Insecticides	Amount/ 100 Gal.	Amount/Gal.
Dormant Oil (2%)	2 gal.	5 tbsp
(3%)	3 gal.	7 ½ tbsp
Malathion 25% WP	3 lb	2 tbsp
Sevin 50% WP	2 lb	2 tbsp
Thiodan 50% WP ¹	2 lb	2 tbsp
Thiodan 3 EC	2/3 qt	1 1/3 tsp
¹ Use only as trunk sprays for peach tree borer.		
Fungicides	Amount/ 100 Gal.	Amount/Gal.
Captan 50% WP	2 lb	2 tbsp
Chlorothalonil	1 lb	1 tbsp
Ferbam 75% WP	2 lb	2 tbsp
Liquid lime-sulfur	12 gal.	2 c
Mancozeb 80% WP	2 lb	2 tbsp
Maneb 80% WP	2 lb	2 tbsp
Myclobutanil (Immunox)	—	½ - 2 oz
Streptomycin	see label	see label
Thiophanate-methyl	6 oz	1 tsp
Wettable Sulfur 95% WP	6 lb	3 tbsp

Harvest Restrictions (Days to Wait after Spraying Until Picking)

Pesticides are poisons, and many backyard fruit growers are concerned about eating fruits that have been sprayed with pesticides.

The U.S. Environmental Protection Agency has established set time periods between the last application of a pesticide (which varies for different pesticides and crops) and harvest to avoid unnecessary residues on ripe fruit. When pesticides are used in accordance with label recommendations, fruit residues are well below the established level allowed for sale. Simply washing fruit before consuming it removes most remaining residue.

There is no problem with excessive pesticide residues on your harvested fruits if:

- Only registered pesticides that list the particular crop to be treated on the label are used.
- Pesticide is used only at the rate recommended on the label.
- Fruit is not picked earlier than the safe preharvest interval given on the label and in Tables 10 and 11.
- Fresh fruit is washed before consumption or processing, even if pesticides are not used.
- Every pesticide label provides harvest restrictions. If your pesticide and its label do not give this information, then it probably is not intended for use on fruit and should not be used.

Dry Formulations

Measurement Equivalents

1 pound = 16 ounces
= 454 grams

1 ounce = 28.4 grams

Sample Calculation for Determining Pesticide Concentration in English Units

1 pound of fungicide per 100 gallons of water
= 16 ounces of fungicide per 100 gallons of water
= 1.6 ounces of fungicide per 10 gallons of water
= 0.16 ounces of fungicide per 1 gallon of water

Sample Calculation for Determining Pesticide Concentration in Metric Units

1 pound of fungicide per 100 gallons of water
= 454 grams of fungicide per 100 gallons of water
= 45.4 grams of fungicide per 10 gallons of water
= 4.5 grams of fungicide per 1 gallon water

Liquid Formulations

Measurement Equivalents

1 gallon = 4 quarts
= 8 pints
= 16 cups
= 128 fluid ounces
= 3785 milliliters

1 quart = 2 pints
= 4 cups
= 32 fluid ounces
= 946 milliliters

1 fluid ounce = 2 tablespoons
= 29.6 milliliters

1 tablespoon = 3 teaspoons
= 14.8 milliliters
= 1/2 fluid ounce

Sample Calculation for Determining Pesticide Concentration for English Units

1 quart of pesticide per 100 gallons of water
= 32 fluid ounces per 100 gallons of water
= 3.2 fluid ounces per 10 gallons of water
= .32 fluid ounces per 1 gallon of water

Sample Calculation for Determining Pesticide Concentration for Metric Units

1 quart of pesticide per 100 gallons of water
= 946 milliliters per 100 gallons of water
= 94.6 milliliters per 10 gallons of water
= 9.4 milliliters per 1 gallon of water

Trade Names	Common Names	Harvest Restrictions/Limitations (Days before Harvest)								
		Apple	Pear	Peach	Cherry	Plum	Blackberry Raspberry	Strawberry	Grape	Blueberry
Agri-strep	streptomycin	50	30	**	**	**	**	**	**	**
Captan	captan	0	*	0	0	0	–**	0	0	0
Carbamate	ferbam	7	7	21	0	–**	40	14	7	40
Immunox	myclobutanil	14	**	0	0	0	**	**	14	**
Ortho Daconil	chlorothalonil	**	**	*	*	*	**	**	**	**
Dithane M-45	mancozeb	77	77	–**	–**	–**	–**	–**	66	–**
Kocide 101 C-O-C-S & others	copper hydroxide, copper oxychloride, fixed copper, tribasic copper sulfate, basic copper sulfate	**	**	**	**	**	**	**	0	**
Sulfur	sulfur	0	0	0	0	0	0	0	0	0
Topsin-M	thiophanate-methyl	0	–**	1	1	1	–**	1	–**	–**

* Limited number of applications allowed or other restrictions apply - REFER TO LABEL DIRECTIONS.
 ** Not registered or recommended.
 NOTE: Check labels for **RE-ENTRY TIMES**. Restrictions in re-entry times may prohibit the use of certain pesticides during harvest.

Trade Names	Common Names	Harvest Restrictions/Limitations (Days before Harvest)								
		Apple	Pear	Peach	Cherry	Plum	Blackberry Raspberry	Strawberry	Grape	Blueberry
Align	azadirachtin	0	0	0	0	0	0	0	0	0
Lorsban	chlorpyrifos	Prebloom	*	14*	14	*	**	21*	35*	**
Malathion	malathion	3	1	7	3	3	1	3	3	8 hrs.
Pyrethrum		1	1	1	1	1	1	1	1	1
Sabadilla		0	0	0	0	0	0	0	0	0
Sevin	carbaryl	3	3	3	3	3	7	7	7	7
Thiodan	endosulfan	21-30*	7*	21-30	21*	7*	**	4*	7	*
Vendex	fenbutatin-oxide	14*	14*	14*	14*	14*	**	1	28	**

* Limited number of applications allowed or other restrictions apply - REFER TO LABEL DIRECTIONS.
 ** Not registered or recommended.
 NOTE: Check labels for **RE-ENTRY TIMES**. Restrictions in re-entry times may prohibit the use of certain pesticides during harvest.

Notes on Pesticide Products Commonly Used by Home Fruit Growers

Even experienced gardeners are sometimes confused by the array of pesticides and special formulations available and the need to select the right materials. Pesticides may be available in one or more formulations.

Wettable Powders. Pesticides used on fruit crops are most commonly sold as wettable powders that are meant to be mixed with water, then sprayed on the crop. If the active ingredient makes up 50 percent of a wettable powder product, it is called a “50W” or “50WP.”

Liquid concentrates are also meant to be mixed with water, then sprayed on the crop. The active ingredient usually ranges from 20 percent to 50 percent of the product.

Dusts are ready to apply as purchased; they are not mixed with water, and they usually contain 1 percent to 10 percent active ingredient.

The following comments will help you select the most appropriate materials for your needs.

Insecticides

Azadirachtin (Align, 3% liquid concentrate) is a botanical insecticide derived from the Neem tree. Azadirachtin, extracted from the seed, is an insect growth regulator and feeding deterrent for pests. It is effective on fruit tree leafrollers, codling moth, Eastern tent caterpillar, peachtree borer, aphids, grape leafhoppers, strawberry rootworm, stink bug and many other insects. It is most effective on the juvenile or larval stages of insects rather than the adult stage. The growth regulator restricts larvae from molting. Results are evident in 3 to 15 days as reduced feeding and death. With repeated applications, insect pressure and plant damage are reduced. For organic certified growers, this chemical has been considered as a “Restricted Use Organic” which allows the use of this pesticide as a final choice alternative. Purchase only enough for one year, since it loses its effectiveness.

Carbaryl (Sevin 50WP, 27% liquid concentrate, 5% dust, 10% dust) is a carbamate insecticide and is effective against Japanese beetle, codling moth, leafrollers, apple maggot, periodical cicada, grape berry moth, oriental fruit moth, flea beetles, spittlebugs, blueberry maggot and many other fruit

insects. Although toxic to aphids, it can actually worsen aphid and mite problems because it is also toxic to their natural enemies. In addition, carbaryl is very toxic to honeybees and its use should be avoided during bloom when bees are active in the planting.

Carbaryl causes apple and pear fruit to drop if used within 21 days of bloom. In some cases, it can result in excessive fruit drop, particularly if a number of sprays are applied. It is often found in all-purpose fruit sprays. Its use should be delayed until four to five weeks after petal fall to avoid thinning. If fruits need to be thinned on a few trees it is best to do it by hand. If there are a number of trees to thin, carbaryl is a readily available thinning agent and works well on a number of cultivars. See ID-92, *Commercial Tree Fruit Spray Guide*, for the specifics on thinning. Too heavy a fruit load can cause tree limbs to break and leads to the production of small fruit and insufficient flower formation for the following year's crop.

Endosulfan (Thiodan 50WP) is a chlorinated hydrocarbon insecticide and is very effective for controlling peach tree borer and the lesser peach tree borer on all stone fruits.

Lindane (Isotox 20% liquid concentrate) is a chlorinated hydrocarbon insecticide and is suggested only as a trunk spray to control peach tree borer.

Malathion (57% liquid concentrate) is an organophosphate insecticide that is relatively safe for general use. It is effective against aphids and other pests. Malathion is not effective where mites are resistant to other organophosphate materials. Its residual effectiveness is generally less than three to four days, so it is especially useful near harvest for pests such as sap beetles, yellowjackets, wasps and bees. Make sure to observe the pre-harvest interval.

Pyrethrum (liquid or WP or dust) is a botanical insecticide extracted from the flowers of a species of *Chrysanthemum*, *Dendranthema grandiflore* from Kenya and Ecuador. This is a natural pyrethrum and should not be confused with the synthetic pyrethroid materials. Pyrethrins give rapid knock-down of caterpillars, beetles, bugs, leafhoppers and leafminers. Pyrethrins leave little residue, degrade quickly and must be sprayed on the insect. Pyrethrins can be harmful to beneficial insects.

Sabadilla (dust) is a botanical insecticide obtained from the seeds of a day lily-like plant native to Central and South America. It is used to control codling moth, aphids and bugs. It has low mammalian toxicity, but can be irritating to the throat and eyes and cause severe sneezing if the applicator is exposed to the floating dust during application. It is harmful to honeybees and breaks down rapidly in sunlight.

Spray oil ("superior," "horticultural," "dormant") is a superior grade of petroleum oil formulated with an emulsifying agent for easy mixing with water. Oil is generally considered an organic treatment. It is used to control pear psylla, mites (European red mite, pear leaf blister mite), and scale insects on tree fruits and is usually applied during the late-dormant to green-tip stage. To avoid severe injury to trees, do not apply when temperatures are likely to fall below 40 degrees F within 24 hours. Thorough coverage of all bark surfaces is very important for oil sprays to be effective.

Microbial Insecticides

Microbial insecticides are derived from beneficial microorganisms such as bacteria, viruses and fungi. These microorgan-

isms cause lethal sickness in pest insects. The most common microbial insecticide is a bacterium, *Bacillus thuringiensis* (Bt). The active ingredient is not the Bt bacterium itself, but a toxin produced by the bacterium that is lethal when ingested by caterpillars, including leafrollers, gypsy moth and codling moth. Because Bt must be consumed to take effect, it is slower acting than other pesticides. Bt is safe not only to humans, but also to beneficial insects such as ladybird beetles, lacewings, predaceous bugs and parasitic wasps.

Insecticidal Soaps

Insecticidal soap is a blend of potassium salts of naturally occurring fatty acids that is more effective for pest control and less phytotoxic than ordinary soaps. However, these soaps have no residual activity and are effective only against insects present at the time of application. There are many home recipes made from dish soap, but dish soap has not been cleared for use on food by the EPA.

Miticides (Acaricides)

Spray oil ("dormant," "horticultural," "superior") is very important for controlling mites on apples and pears. See comments on oil under "Insecticides."

Special note concerning mite build-ups

Sevin and sulfur are ineffective against pest mites, but are highly toxic to predator mites. As a result, mite problems may develop on trees after the use of these materials.

Fungicides

Captan (Orthocide 50WP) is a broad-spectrum fungicide for use on most fruit crops and many vegetable and ornamental plants. Captan controls most diseases listed under thiophanate-methyl and is often used in combination with thiophanate-methyl. It also helps control peach leaf curl. It does not control powdery mildew or rust diseases. Because of incompatibility and injury problems, Captan should not be used in combination with either sulfur or spray oil. Captan causes severe leaf injury on Japanese-type and Stanley plums if used before July.

Chlorothalonil (Ortho Daconil 2787) is a broad spectrum protectant effective against peach leaf curl and fruit diseases of peach, plum and cherry. It can only be used early in the growing season for fruit rot control.

Ferbam (Carbamate 76WP), a black, dry wettable powder, is a broad-spectrum fungicide, with limited use on fruit crops. It is very effective for control of cedar apple rust and as a dormant spray for peach leaf curl. It is more effective than thiophanate-methyl or Captan for control of grape black rot. Ferbam leaves a black, unsightly residue on treated plant surfaces; therefore, its use should be restricted to early in the growing season.

Mancozeb (Dithane M-45, F-45, DF) is a yellow, dry wettable powder or flowable fungicide with broad-spectrum protectant activity. It is good for control of scab and rust diseases of apple, and for control of grape black rot, leaf spot, and downy mildew.

Myclobutanil (Immunox, Nova) is a locally systemic fungicide effective against apple scab and cedar apple rusts. It can also be used for managing brown rot of peaches and other stone fruits and for black rot and anthracnose of grape. It is very effective against powdery mildew diseases of apples, stone fruits, and grapes. It is best used in combination with

captan or mancozeb, and should not be used on apples past first cover.

Thiophanate-Methyl (3336, Topsin-M) is a liquid systemic fungicide effective against powdery mildew and other diseases of apple and stone fruits. On apples, it controls scab, powdery mildew, blotches and fruit rots, but not rust diseases. On stone fruits, it controls brown rot, scab, black knot of plums and cherry leaf spot. Use in combination with Captan or other labeled fungicide when possible.

Copper Fungicides

The original material used was copper sulfate (also known as blue vitriol or blue stone). Although copper controls fungi and bacteria, it also injures plants. When this material was combined with lime in the French vineyards, the combination became known as Bordeaux mixture.

Bordeaux Mixture is a mixture of copper sulfate and hydrated lime in water. It has long residual action and has been used to control many diseases, including peach leaf curl, fire blight and scab of apple and black rot, downy mildew and powdery mildew of grape. It is available as a pre-mixed dry wettable powder or in liquid form.

Fixed Copper Fungicides. Following the discovery and use of Bordeaux mixture, several relatively insoluble copper compounds or fixed coppers were developed. Fixed copper formulations are less injurious to plant tissues than Bordeaux mixture, but their use is still limited because of their potential to injure plants and lack of compatibility with other pesticides. Some more common formulations of fixed copper include C-O-C-S (copper oxychloride sulfate), Kocide 101 (copper hydroxide), Tribasic Copper Sulfate, and basic copper sulfate.

Sulfur Fungicides

Liquid Lime Sulfur. Liquid lime sulfur fungicide can be used as a dormant spray on peaches for peach leaf curl, on plums for black knot, and on raspberries and blackberries for cane blight, spur blight and anthracnose. It should be used only when plants are dormant. It can cause severe damage if applied after green foliage appears.

Dry Wettable Sulfurs. Dry wettable sulfurs are available under many trade names. Microfine wettable sulfurs are usually much less injurious to foliage and fruit than liquid lime sulfur, but their use during hot weather (above 85 degrees F) may result in leaf burning and fruit russetting. Dry wettable sulfurs are very effective for control of powdery mildew on most fruit crops. **Sulfur is very toxic to foliage of certain grape varieties (mainly American grapes) including 'Concord' and the French hybrid grapes 'Chancellor,' 'DeChaunac' and 'Foch.'** Sulfur is relatively safe on most other varieties. Applications after fruit begins to ripen may pose problems during fermentation if grapes are intended for wine making.

Sulfur is lethal to some beneficial insects, spiders and mites. These beneficial insects are natural predators of harmful insects and mites that affect fruit crops. Killing beneficial insects may increase certain pest problems, especially mites.

Sulfur provides good control of brown rot and scab of stone fruits. Sulfur is only moderately effective against apple scab. For good control, sulfur applications must be perfectly timed ahead of scab infection periods.

Bactericides

Copper Materials. On apples and pears where fire blight was present the previous season, an application of Bordeaux mixture or fixed copper materials may be beneficial. Spray thoroughly in spring just before green tissue appears. Where bacterial spot has been a problem on peaches, an application of fixed copper just as leaves begin to fall in autumn helps reduce damage from this disease the next season. Copper materials applied during the growing season can damage apple leaves and fruit and cause complete defoliation of peaches and other stone fruits.

Streptomycin. Streptomycin is an antibiotic of limited use, but is important for apple and pear varieties with a history of fire blight problems. Although streptomycin is commonly used in commercial apple orchards and is not a restricted-use pesticide, it may be difficult for homeowners to obtain. In addition, timing and number of required sprays are critical to achieve effective control and to avoid problems with resistance. For all these reasons, growers should select varieties with some natural resistance to fire blight and employ cultural practices that reduce risk of infection.

Multipurpose or All-Purpose Fruit Sprays

Insecticides and fungicides are often needed at the same time on a plant to protect it against attack by both insects and diseases. All-purpose fruit sprays are available. These contain a mixture of insecticides and fungicides to give adequate control for most common problems and are easy to use. If a mixture containing Sevin is used, do not apply it until 21 days after petal fall on apples and pears. Sevin thins or causes fruit to drop if applied before this date. Sevin may also cause buildup of pest mites.

These commercially prepared mixtures are convenient to use and reduce the number of pesticides the homeowner must store safely. Other considerations include:

- None of these multipurpose mixtures controls all insects and diseases you are likely to encounter. Not realizing this, many users tend to apply mixtures more frequently and at higher rates trying, in vain, to control some pests.
- Multipurpose mixtures lack flexibility so that when only a fungicide is required (e.g., during bloom), an insecticide also is applied even though it is not needed and could be damaging.
- Most multipurpose mixtures are more expensive than those you prepare yourself.

You can mix all-purpose fruit spray with additional pesticides, but there are some limitations with which you should become familiar. If you are in doubt about spray combination injury, check the label. Apply the mixture to a small portion of the tree and wait to see what happens. Spray burn should show up in 24 to 48 hours. If burn results, do not apply the combination. If you plan to mix your own combination, use wettable powders to reduce plant injury.

Where to Purchase Pesticides

Most garden centers or stores with yard and garden departments where pesticides are sold have commonly recommended pesticides. Many agricultural supply or farmer co-op centers also carry them.

Organic Pesticides

Organic culture of fruit crops is a rewarding and often a very challenging undertaking. **Organic production is not a passive means of pest control, but a production system in which a variety of cultural practices, organic pesticide controls and beneficial insects are used.** Considerable care must be taken in determining the planting location, layout, pest control and cultural practices. The correct cultural practices are extremely important to follow when growing fruit using organic techniques. See the recommendations for the specific fruit crop of interest throughout this publication. Picture perfect fruit may not be consistently possible with the extremes of climate and the pests encountered in Kentucky. Follow the proper management techniques, scout for problems before economic thresholds are met and keep informed of the latest cultural information. Generally the small-fruit crops, excluding grapes, are much easier to grow organically than tree fruit crops. Most botanical insecticides have a broad spectrum of activity and users should be aware that many of these materials are toxic to beneficial insects and mites as well as pest insects. Some botanical insecticides are as toxic as some synthetic insecticides and should be handled accordingly.

If fruit is to be sold as organically certified in Kentucky, contact the Kentucky Department of Agriculture, Division of Markets, Frankfort, KY, phone 502/564-4696, for the certification requirements. Only certain pest control materials may be used for insect and disease control. Some of these include the insecticides pyrethrin (without piperonyl butoxide) Sabadilla; insecticidal soaps; Bt species; horticultural oils (non-synthetic); herbal extracts and oils, diatomaceous earth and the fungicides; copper and sulfur (only certain preparations of these are acceptable).

Nearly all of these materials can be used up to the day of harvest.

Table 12 on page 17 contains a list of organic insecticides and practices that people have tried for insect and mite control on the various fruit crops. You may want to experiment with some of these insecticides and practices on small plantings. **Many of these insecticides and practices are not necessarily as effective as other recommended control measures, and research data to support the use of some of these is lacking.**

One method of dealing with pests sustainably is through the introduction of beneficial insects and predators of the insect pests (Table 13). Beneficial insects do not ‘cure’ the pest problem in a quick fashion as do pesticidal means. They take time and work on reducing the overall population of the pest to economically tolerable levels if the instructions set by the insectary are followed. First, buy your insects through a reputable dealer. The beneficials should be included as part of an integrated pest control system that includes fertilizing, proper cultural practices as outlined within this publication, the use of resistant varieties of fruit when possible, proper irrigation, and timely monitoring of pests.

Using beneficials is most effective when pests are already present in your planting at low or moderate levels. The beneficial insects need an immediate food source to start with. To maintain a constant level of beneficials, there must always be some food source available. So the goal of this type of pest management scheme is not the total eradication of the pests, only the managed balance between beneficial and pest.

Target Insect	Beneficial Insect Control
All species of soft-bodied insect, mites, eggs	Green Lacewing <i>Chrysoperla carnea</i> , <i>C. rufilabris</i>
Eggs of most species of moths and butterflies	Parasitic wasps <i>Trichogramma minutum</i> , <i>T. platneri</i> , <i>T. pretiosum</i>
Two-spotted spider mite	Predatory mites <i>Amblyseius cucumeris</i> , <i>A. fallacis</i>
Various mites	Predatory mites <i>Neoseiulus californicus</i> , <i>Galendromus occidentalis</i> , <i>Phytoseiulus persimilis</i> , <i>Mesoseiulus longipes</i>
Strawberry crown borer	Predatory nematodes <i>Steinerema carpocapsae</i>
Strawberry root weevil	Predatory nematodes <i>Steinerema carpocapsae</i>
Strawberry rootworm	Predatory nematodes <i>Steinerema carpocapsae</i>
Walnut husk fly	Predatory nematodes <i>Steinerema carpocapsae</i>
Raspberry cane moth	Predatory nematodes <i>Steinerema carpocapsae</i>
Pear Psylla	Predatory bug <i>Deraeorcoris brevis</i>
Thrips and mites	<i>Iphiseius degenerans</i>
Leaf miners	Parasitic wasp <i>Diglyphusisea isea</i>

Monitoring Insect Pests

Codling Moth Trap—Pheromone lure with sticky trap. Set trap out two weeks before buds break. If later in season, traps would be effective against second or third generation infestations. Lures need to be replaced every 4 weeks. For larger sized trees, at least 2 traps would be necessary. (See ENTFACT 203)

Oriental Fruit Moth—Pheromone lure monitoring trap (See ENTFACT 212)

Peachtree Borer—Pheromone lure monitoring trap (See ENTFACT 200)

Lesser Peachtree Borer—Pheromone lure monitoring trap (See ENTFACT 200)

Strawberry Crown Borer—Pheromone lure monitoring trap (New and guidelines have not been established)

San Jose Scale—Pheromone lure monitoring trap (See ENTFACT 204)

Monitoring Traps:

Where to place

Place traps in the southeast corner of the tree canopy, 5 to 7 feet above the ground for tree crops. Traps should be placed on the perimeter of a planting facing the predominant wind direction or bordering wooded areas. In larger plantings, traps could be positioned at two corners and one at the interior.

When to start

Traps should be placed into the planting at the pink stage in tree crops. An early start allows the tracking of the first generation from the very beginning. Differentiate between older and younger trees.

Monitoring





















Traps should be checked weekly except during peak emergence times when they are checked daily. Record the results. Further information on apple integrated pest management is available in IPM-9, *Kentucky Backyard Integrated Pest Management Manual*.






















Table 12. Organic Insect and Mite Control for Fruit

Insects	Fruit Type ¹	Pyrethrum	Sabadilla	Align	Entrust	Surround Crop Protectant	Insecticidal Soaps	Bt Types ³	Dormant Oils ⁴	Citrus Peel Oils ⁵	Garlic Oil ⁶	Sticky Traps ⁷	Barriers ⁸	D.E. ⁹	Cultural Practices/ Comments ¹⁰
Aphids	All	a ²	c	b			a			a		a, yellow trap for monitoring only sticky band around trunks ¹¹			
Apple Tree Borers	A														
Blackberry Crown Borer	B, R														
Codling Moth	A, Pr	a	a	b	a			a					a		pick up and destroy fallen fruit
Fall Webworm	All tree fruit			b	a										submerge tents in soapy water
Flea Beetles	G	a	b							a				a	
Japanese Beetle	All												fine netting for some crops		
Leafhopper Grape	G			b			a								
Leafroller	A, G, Pl, Pr	a		b	a			a							
Oriental Fruit Moth	A, P	b	a	a											
Peachtree Borer	C, P, Pl			b								sticky bands at tree base ¹¹			kill borers with wire or knife carefully
Pear Psylla	Pr	a				a								a	
Plum Curculio	A, Blu, C, P, Pl, Pr					a									
Raspberry Cane Borer	B, R														cut out infected canes and burn
Red-Necked Cane Borer	B, R														cut out swollen areas in canes and burn
Strawberry Rootworm	S			b				a						b	
Scale	Ch, P, Pl								dormant oil						
San Jose Scale	A, C, P, Pl, Pr														same as above and use double-sided tape to monitor crawlers
Spider Mite	All	a				a		a		a					alternate spray type
Strawberry Crown Borer	S														
Strawberry Root Weevil	S	a		a											
Stinkbug	A, B, P, Pr, R	b	b	b											
Tarnished Plant Bug	A, B, P, Pr, R	a	a												a
Tent Caterpillar	A, C, P, Pl, Pr			b	a			a							submerge tents in soapy water

1 Fruit type, A=apple, B=blackberry, Blu=blueberry, C=cherry, G=grape, P=peach, Pl=plum, Pr=pear, R=Raspberry, S=Strawberry
 2 a = the primary control method, b= secondary control method, c= somewhat effective method
 3 Bt types are different strains of the Bt bacteria (*Bacillus thuringiensis*) that attack certain groups of insect larvae. Bt var kurstaki controls some moth and butterfly larvae and Bt var tenebrionis controls some beetle larvae.
 4 Dormant oils are specially refined oils that are meant to smother the insect pests while not injuring the plants. Labels must be carefully read to be sure that no other chemicals have been added and that these oils comply with Organic Certification Standards.
 5 Citrus Oil Spray is a very successful contact insecticide that immobilizes insects in a few minutes. It is nonselective and can be harmful to some beneficial insects. Chop orange peels and place in a pan with just enough water to cover. Simmer for 5 minutes. Drain off the liquid after cooling and use.
 6 Garlic Oil Spray must be applied as soon as it is made. It will kill all types of insects including beneficials so use sparingly. Use 3-5 tsp. of garlic cloves chopped. Add 3-5 tsp. mineral oil and soak for 24 hours. Mix 1 pt. with tsp. insecticidal soap then add slowly to the garlic mixture. Stir and strain. Use 2 tbsp. in 1 pint of water for spray.
 7 Sticky Traps attract insects with color and catch them on sticky resin. While these are available commercially, they can be made at home. Use wooden squares, plastic sheets, or heavy waxed cardboard sprayed or painted with glossy yellow, red, or white colors depending on the pest. Spread on mineral oil or petroleum jelly mixed with kitchen detergent. Hang traps near the fruit crop.
 8 Barriers can be netting or finely woven poly cloth which lightly covers plants that physically separates plants from the insects. All sides must be securely fastened to the ground. It will raise the temperature and humidity under the cover so care must be taken to avoid overheating the plants on hot days. It can be quite expensive to purchase but the high quality material will last for years.
 9 D.E. Diatomaceous earth is the powdered remains of fossilized diatoms. The powder has extremely small but sharp protrusions that severely injure insects when they crawl over it. Harmless to humans and animals. Reapply after each rain.
 10 Cultural Practices is a broad category. Most often, it includes the removal of overwintering plant debris and fruit drops from the field or garden plot. This deprives insects of early emergence and establishment in the spring. Many cultural practices can be done during the growing season. Ideally, the best control for organic growers is prevention. The best prevention comes from the maintenance of a living healthy soil that keeps plants growing at an optimum level. Stressed plants are much more prone to infestation and infection than healthy ones. Proper humus, nutrient, and moisture levels are essential for maintaining a balanced soil environment.
 11 Placing sticky bands around the trunk involves wrapping burlap around the trunk, burying it several inches below ground and covering it with tanglefoot.
 Wormwood (*Artemisia absinthium*) extracts from the leaves are used as a soil drench or spray for repelling slugs and snails. Steep leaves in boiling water, let cool and use immediately.
 Liquid Seaweed Extract will disrupt the life cycle of many insects when used diluted with water. Mix with insecticidal soap for foliar spray that offers a broad range of pest protection.

Floral Development Stages for Fruit Crops and Critical Temperatures for Flower Bud Kill

	Apple	Pear	Peach	Tart Cherry	Plum and Prune
Stage 1	<p>Dormant</p> 	<p>Dormant</p> 	<p>Dormant</p> 	<p>Dormant</p> 	<p>Dormant</p> 
Stage 2	<p>Silver tip</p>  <p>10% kill 15° F 90% kill 2° F</p>	<p>Swollen bud</p>  <p>10% kill 15° F 90% kill 1° F</p>	<p>Swollen bud</p>  <p>10% kill 18° F 90% kill 2° F</p>	<p>Bud burst</p>  <p>10% kill 17° F 90% kill 5° F</p>	<p>Swollen bud</p>  <p>10% kill 14° F 90% kill 1° F</p>
Stage 3	<p>Green tip</p>  <p>10% kill 18° F 90% kill 10° F</p>	<p>Bud burst</p>  <p>10% kill 20° F 90% kill 7° F</p>	<p>Half-inch green</p>  <p>10% kill 23° F 90% kill 5° F</p>	<p>Green tip</p>  <p>10% kill 25° F 90% kill 14° F</p>	<p>Bud burst</p>  <p>10% kill 18° F 90% kill 3° F</p>
Stage 4	<p>Half-inch green</p>  <p>10% kill 23° F 90% kill 15° F</p>	<p>Green cluster</p>  <p>10% kill 26° F 90% kill 15° F</p>	<p>Pink</p>  <p>10% kill 25° F 90% kill 18° F</p>	<p>Tight cluster</p>  <p>10% kill 26° F 90% kill 17° F</p>	<p>Green cluster</p>  <p>10% kill 26° F 90% kill 16° F</p>

<p>Stage 5</p>	<p>Tight cluster</p>  <p>10% kill 27° F 90% kill 21° F</p>	<p>White bud</p>  <p>10% kill 26° F 90% kill 22° F</p>	<p>Bloom</p>  <p>10% kill 27° F 90% kill 24° F</p>	<p>Swollen bud</p>  <p>10% kill 27° F 90% kill 24° F</p>	<p>White bud</p>  <p>10% kill 26° F 90% kill 21° F</p>
<p>Stage 6</p>	<p>Pink</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Bloom</p>  <p>10% kill 28° F 90% kill 23° F</p>	<p>Petal fall</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Bloom</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Bloom</p>  <p>10% kill 27° F 90% kill 23° F</p>
<p>Stage 7</p>	<p>Bloom</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Petal fall</p>  <p>10% kill 28° F 90% kill 24° F</p>	<p>Fruit set- shucks on</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Petal fall</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Petal fall</p>  <p>10% kill 28° F 90% kill 23° F</p>
<p>Stage 8</p>	<p>Petal fall</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Fruit set</p>  <p>10% kill 28° F 90% kill 24° F</p>	<p>Fruit set- shucks off</p> 	<p>Fruit set</p>  <p>10% kill 28° F 90% kill 25° F</p>	<p>Fruit set</p> 
<p>Stage 9</p>	<p>Fruit set</p>  <p>10% kill 28° F 90% kill 25° F</p>				

Mention or display of a trademark, proprietary product or firm in text or figures does not constitute an endorsement and does not imply approval to the exclusion of other suitable products or firms.

Educational programs of Kentucky Cooperative Extension serve all people regardless of race, color, age, sex, religion, disability, or national origin. Issued in furtherance of Cooperative Extension work, Acts of May 8 and June 30, 1914, in cooperation with the U.S. Department of Agriculture, M. Scott Smith, Director of Cooperative Extension Service, University of Kentucky College of Agriculture, Lexington, and Kentucky State University, Frankfort. Copyright © 2005 for materials developed by University of Kentucky Cooperative Extension. This publication may be reproduced in portions or its entirety for educational or nonprofit purposes only. Permitted users shall give credit to the author(s) and include this copyright notice. Publications are also available on the World Wide Web at www.ca.uky.edu.

Revised 4-2005.